

The Art and Science of Improving Lives Prosthetic and Orthotic Services

Donning/Doffing of Double Wall Transfemoral EPN Socket System

- 1) Residual Limb Inspection
 - Inspect residual limb each day prior donning your prosthesis
 - * Use small hand held mirror to view distal end
- 2) Apply Skin lubricant
 - Anatomical Designs Inc. recommends A&D Ointment, Monistat Soothing Care Chafing Relief Powder Gel
- 3) Donning Silicone Liner
 - Invert silicone Liner
 - Place against distal end of residual limb
 - Roll silicone liner up your residual limb *Burp silicone liner*
 - Inspect to ensure liner has no air trapped between it and your residual limb
- 4) Place Air Wick
 - Nylon
 - Appropriate ply of sock
- 5) Donning Distal Cup
 - Sealing sleeve should be inverted down the outside of distal cup
 - Place distal cup on residual limb over silicone liner and air wick
 - Position distal cup in appropriate position
 - Reflex sealing sleeve up over distal cup, air wick, and silicone liner

- 6) Charge Vacuum Camber
 - Charge chamber to value established by your Prosthetist (Have lines to fill in with vac values)
- 7) Don Retainer
 - Slide retainer on and secure side pin, lanyard, or distal pin
 - •On side pin insure that blue button snaps back into closed position

1-800-418-0313

www.anatomicaldesignsinc.com